

HARBS PLUS

Phytogenic Anti-Viral Drugs for Poultry and Dairy

Composition:

Andogilio, Afvogilio, Pengiouactia, Subulla, Coriandrum, Caryophyllue, Zerum, Longissima, Bioactive peptides and Amino Acids.

Indication:

For prevention and treatment of New castle disease, Infectious bursal disease, Infectious bronchitis, Infectious laryngotracheitis, E.D.S, Fowl Pox, Marek's, Duck plague, Poultry fever, Feed intake decreased, Coughing, swelling of the eyelids, green or off-white feces, laying rate drop, poor quality egg shell and so on. Animal viral diseases like Foot and Mouth disease (FMD), P.P.R., anorexia and so on. Mixed with antibiotics in the treatment of mixed infections.

Lifting immunosuppression:

Fungi poisoning, poultry's bursal disease, coccidiosis caused immunosuppression.

Features:

- To inhibit and destroy the viruses.
- Enhancing body's immune and cellular immune functions.
- Reducing morbidity and mortality quickly.
- Adjuvant treatment of mixed infections.

Dosage and Administration:

Viral Diseases (Including mixed infections):

Poultry: Add 100 gm HARBS Plus into 300-500 Ltr. Of drinking water or use 0.05 gm per kg body weight orally via drinking water twice daily for 5-7 days. Prevention dosage halved.

Livestock: 0.05-0.1 gm per kg of body weight twice daily for 7-15 days. Prevention dosage halved.

Severe case: Three times of the normal amount by feed or drinking water and next days are repeat.

Note: Limited water 1-2 hours before mixed drinks for using the drug. To enhance antibacterial effect if combining with antibiotics.



Precaution:

Not for human, animal use only. Keep out of reach of children. Store in a cool and dry place, protect from direct sunlight and moisture.

Pack Size: 100 gm mono cartoon.







